State of New Jorsey DEPARTMENT OF EDUCATION

Sudden Cardiac Death Pamphlet Sign-Off Sheet

Name of School District:	7
Name of Local School:	
	9.8.
I/We acknowledge that we received and review	wed the Sudden Cardlac Death in Young Athletes pamphiet.
39	
e.	
Student 91 gnature:	
Parent or Guardian	
Date:	

Website Resources

- Sudden Death in Athletes www.cardiachealth.org/sodden-death-thathletes
- Hypertrophic Cardlomyopathy Association www.4licm.org
- Annerican Heart Association www.heart.org

Collaborating Agencies:

American Academy of Podlatrics American Academy of Podlatrics New Jersey Chapter 3036 Quakerbridge Road, Suite 108 Hamilton, NJ 08619 (p) 609-842-0014 (f) 609-842-0015 www.aapn).org



American Heart Association 1 Union Street, Suite 301 Robbinsville, NJ, 08691 (p) 609-208-0020



www.hearLorg New Jersey Department of Education PO Box 500

(p) 609-292-5935 www.statenj.us/education/




Trenton, NJ 08625-0360 (p) 609-292-7837 www.state.nj.us/health


Lead Author: American Academy of Pediatrics, New Jersey Chapter

Written by: Initial draft by Sushma Raman Hebbar, MD & Stephen G. Rice, MD PhD

Additional Reviewers: NJ Department of Eduration, NJ Department of Health and Senior Services, American Heart Association/New Jersey Chapter, NJ Academy of Family Practice, Pediatric Cardiologists, New Jersey State School Nurses


Revised 2014: Christene DeWitt-Parker, MSN, CSN, RN; Lakota Kruse; MD, MPH; Susan Martz, EdMc Stephen G, Rice, MO; Jeffey Rosenberg, MD, Leuis Teichholz, MO; Purry Weinstock, MD











udden death in young athlete between the ages of 10 and 19 is very rare. What, if anything, can be done to prevent this kind tragedy?

Sudden cardiac death is the result of an unexpected failure of proper heart function, usually (about 60% of the time) during or immediately after exercise without trauma. Since the heart stops pumping adequately, the athlete quickly collapses, loses consciousness, and ultimately dies unless normal heart rhythm Is restored using an automated external defibrillator (AED).

Sudden cardlac death in young athletes is very rare. About 100 such deaths are reported in the United States per year The chance of sudden death occurring to any individual high school athlete about one in 200,000 per year.

Sudden cardiac death is more common: In males than in females: in football and basketball than in other sports: and in African-Americans than In other races and ethnic groups.

Research suggests that the main cause is a loss of proper heart rhythm, causing the heart to quiver instead of pumping blood to the brain and body. This is called ventricular fibrillation (ven-TRICK-you-lar fibroo-LAY-shun). The problem is usually caused by one of several cardlovascular abnormalities and electrical diseases of the heart that go

unnoticed in healthy-appearing athletes.

The most common cause of sudden death in an athleta is hypertrophic cardiomyopathy (ht-per-TRO-fic CAR- dee-ol-my-OP-a-thee) also called HCM, HCM is a disease of the heart, with abnormal thickening of the heart muscle, which can cause serious heart rhythm problems and blockages to blood flow. This genetic disease runs in familles and usually develops gradually over many years.

The second most likely cause is congenital (con-JEN-It-all fle, present from birth) abnormalities of the coronary

arteries. This means that these blood vessels are connected to the main blood vessel of the heart in an abnormal way. This differs from blockages that may occur when people get older (commonly called coronary artery disease," which may lead to a heart

SUDDEN CARDIAC DEATH IN YOUNG ATHLETES

Other diseases of the heart that can lead to sudden death in young people include:

- Myocarditis (my-oh-car-DIE-tis), an acute inflammation of the heart muscle (usually due to a virus).
- Dilated cardiomyopathy, an enlargement of the heart for unknown reasons.
- Long QT syndrome and other electrical abnormallties of the heart which cause abnormal fast heart rhythms that can also run in familles.
- Marfan syndrome, an inherited disorder that affects heart valves, walls of major arteries, eyes and the skeleton. It is generally seen in unusually tall athletes, especially if being tall is not common in other family members.

Wel-Are there warning signs to watch for?

In more than a third of these sudden cardiac deaths, there were warning signs that were not reported or taken seriously. Warning signs are:

- Fainting, a seizure or convulsions during physical activity;
- Fainting or a seizure from emotional excitement, emotional distress or being startled;
- Dizziness or lightheadedness, especially during excition;
- Chest pains, at rest or during exertion;

- Palpitations awareness of the heart beating unusually (skipping, irregular of extra beats) during athletics or during codown periods after athletic participation;
- Fatigue or tiring more quickly than peers, or
- Being unable to keep up with friends due to shortness of breath.

What are the current recommendations:

New Jarroy requires all rchool athletes to be examined by their primary care physician ("medical home") or school physician at least once per year. The New Jersey Department of Education requires use of the specific Annual Athletic Pie-Participation Physical Examination Form.

This process begins with the parents and student-athletes answering questions about symptoms during exercise (such as chest pain, dizziness, fainting, palpitations or shortness of breath); and guestions about family health history.

family health history.

The primary healthcare provider needs to know if any family member died suddenly during physical activity or during a seizure. They also need to know if anyone in the family under the age of 50 had an unexplained sudden death such as drowning or cal accidents. This information must be provided annually for each exam because it is so essential to identify those at risk for sudden cardiac death.

The required physical exam includes measurement of blood pressure and a careful listening examination of the heart, especially for murmurs and rhythm abnormalities. If there are no warning signs reported on the health history and no abnormalities discovered on exam, no further evaluation or testing is recommended.

When should a student at hiete see a heart specialist?

If the primary healthcare provider or school physician has concerns, a referral to a child heart specialist, a pediatric cardiologist, is recommended. This specialist will perform a more thorough evaluation, including an electrocardiogram (ECG), which is a graph of the electrical activity of the heart. An echocardiogram, which is an ultrasound test to allow for direct visualization of the heart structure, will likely also be done. The specialist may also order a treadmill exercise test and a monitor to enable a longer recording of the heart rhythm. None of the testing is invasive or uncomfortable.

Can audden cardiac death be prevented Just through proper screening?

A proper evaluation should find most, but not all, conditions that would cause sudden death in the athlete. This is because some diseases are difficult to uncover and may only develop later in life. Others can develop following a normal screening evaluation, such as an infection of the heart muscle from a virus.

This is why scenning evaluations and a review of the family health history need to be performed on a yearth basis by the athlete's primary healthcare provider. With proper screening and evaluation, most cases can be identified and prevented.

Why have an AED on site during spore events?

The only effective treatment for ventricular fibrillation is immediate use of an automated external defibrillator (AED). An AED can restore the heart back into a normal rhythm. An AEU is also ille-saving for ventricular fibrillation caused by a blow to the chest over the heart (commote Cordis).

the chest over the heart (commot for cords):

Effective September 11 2014, the New Jersey
Department of Education requires that allpublic and nonpublic schools grades K
through 12 shalt

- Have an AED available at every sports event (three minutes total time to reach and return with the AED);
- Have adequate personnel who are trained in AED the present at practices and games;
- Have, coaches and all letic trainers trained in basic life support techniques (CPR); and
- Call 911 immediately while someone is retrieving the AED.